

Theoretical Question 2

Rising Balloon

A rubber balloon filled with helium gas goes up high into the sky where the pressure and temperature decrease with height. In the following questions, assume that the shape of the balloon remains spherical regardless of the payload, and neglect the payload volume. Also assume that the temperature of the helium gas inside of the balloon is always the same as that of the ambient air, and treat all gases as ideal gases. The universal gas constant is $R=8.31 \text{ J/mol}\cdot\text{K}$ and the molar masses of helium and air are $M_H = 4.00 \times 10^{-3} \text{ kg/mol}$ and $M_A = 28.9 \times 10^{-3} \text{ kg/mol}$, respectively. The gravitational acceleration is $g = 9.8 \text{ m/s}^2$.

[Part A]

(a) [1.5 points] Let the pressure of the ambient air be P and the temperature be T . The pressure inside of the balloon is higher than that of outside due to the surface tension of the balloon. The balloon contains n moles of helium gas and the pressure inside is $P + \Delta P$. Find the buoyant force F_B acting on the balloon as a function of P and ΔP .

(b) [2 points] On a particular summer day in Korea, the air temperature T at the height z from the sea level was found to be $T(z) = T_0(1 - z/z_0)$ in the range of $0 < z < 15$ km with $z_0 = 49$ km and $T_0 = 303$ K. The pressure and density at the sea level were $P_0 = 1.0 \text{ atm} = 1.01 \times 10^5 \text{ Pa}$ and $\rho_0 = 1.16 \text{ kg/m}^3$, respectively. For this height range, the pressure takes the form

$$P(z) = P_0(1 - z/z_0)^\eta . \quad (2.1)$$

Express η in terms of z_0 , ρ_0 , P_0 , and g , and find its numerical value to the *two* significant digits. Treat the gravitational acceleration as a constant, independent of height.

[Part B]

When a rubber balloon of spherical shape with un-stretched radius r_0 is inflated to a sphere of radius r ($\geq r_0$), the balloon surface contains extra elastic energy due to the stretching. In a simplistic theory, the elastic energy at constant temperature T can be expressed by

$$U = 4\pi r_0^2 \kappa RT \left(2\lambda^2 + \frac{1}{\lambda^4} - 3 \right) \quad (2.2)$$

where $\lambda \equiv r/r_0$ (≥ 1) is the size-inflation ratio and κ is a constant in units of mol/m².

(c) [2 points] Express ΔP in terms of parameters given in Eq. (2.2), and sketch ΔP as a function of $\lambda = r/r_0$.

(d) [1.5 points] The constant κ can be determined from the amount of the gas needed to inflate the balloon. At $T_0 = 303$ K and $P_0 = 1.0$ atm = 1.01×10^5 Pa, an un-stretched balloon ($\lambda = 1$) contains $n_0 = 12.5$ moles of helium. It takes $n = 3.6 n_0 = 45$ moles in total to inflate the balloon to $\lambda = 1.5$ at the same T_0 and P_0 . Express the balloon parameter a , defined as $a = \kappa/\kappa_0$, in terms of n , n_0 , and λ , where $\kappa_0 \equiv \frac{r_0 P_0}{4RT_0}$. Evaluate a to the two significant digits.

[Part C]

A balloon is prepared as in (d) at the sea level (inflated to $\lambda = 1.5$ with $n = 3.6 n_0 = 45$ moles of helium gas at $T_0 = 303$ K and $P_0 = 1$ atm = 1.01×10^5 Pa). The total mass including gas, balloon itself, and other payloads is $M_T = 1.12$ kg. Now let the balloon rise from the sea level.

(e) [3 points] Suppose that the balloon eventually stops at the height z_f where the buoyant force balances the total weight. Find z_f and the inflation ratio λ_f at that

height. Give the answers in two significant digits. Assume there are no drift effect and no gas leakage during the upward flight.